



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MINOR NOTICES.

Pflanzenfamilien.⁴—Part 226 contains the completion of the Neckeraceae and the beginning of the Lembophylaceae by V. F. BROTHERUS. The second part, of the second supplement has also appeared, including the literature of 1899-1904 in reference to dicotyledons up to the beginning of Euphorbiaceae.—J. M. C.

NOTES FOR STUDENTS.

Archegoniatenstudien.—The tenth of GOEBEL's series by this title is for size almost a book in itself. It is made up of over twenty papers on morphology and biology of mosses and liverworts, varying from a page or less, embodying a brief note on the water adaptation in the form and position of the leaves in Orthorhynchium, to a paper of forty-odd pages on Dawsonia and its allies, and a like one on marsupiferous Jungermanniales. In great part these papers were written several years ago, and some of the researches have been epitomized in GOEBEL's *Organographie*, but they have not been published *in extenso* until now, on account of other work.

Together they form a most important contribution to our knowledge of the bryophytes—a contribution too full of details to report fully. At various places it has a tinge of the polemic, for the author has to clear away many errors, and he takes occasion to rebuke one and another for shortcomings. Much space is devoted to speculations, which are confessedly unsupported by investigation because material or time was lacking. Such speculations, if put briefly, may be suggestive as a guide to future investigations; but they appear to be indulged in as a basis for future claims to priority, if we may judge from some citations of earlier ones in these pages. Even a scientific man is rarely without a prejudice in favor of his own hypotheses. Thus the author guessed (*Organographie* 346) about the development of the multicellular "spores" of Dicnemon: "Am wahrscheinlichsten ist es dass sich aus den gekeimten Sporen ein Fadenprotonema bildet, etwa aus den Brutknospen von Tetraxis." Now he declares "dass die früher geäusserte Vermutung richtig war." But the "protonemal filaments" function "der Hauptsache nach als Rhizoiden;" rhizoids arise also from the surface; and the apical cell of the stem arises not as a branch of one of these "protonemal filaments" but almost immediately from a marginal cell of the "spore." It is difficult to see how the earlier guess can possibly be justified by these observations. Certainly the resemblance to the behavior of the gemmae of Tetraxis is rather remote. Other like instances might be cited; sometimes the guess was right, sometimes not; and that is likely to be the case with these new ones. But the observations are abundant, and the author's keen discrimination and clear presentation throw light upon many obscure points.

Dawsonia is held to be the primitive form of the Polytrichum line by reason of the limited differentiation of tissues in the axis of the gametophyte, and espe-

⁴ ENGLER, A., und PRANTL, K., Die natürlichen Pflanzenfamilien. Lieferung 226. Leipzig: Wilhelm Engelmann. 1906.